

Updated October 30, 2024

Homework problems for AMAT 540A (Topology I), Fall 2024. Over the course of the semester I'll add problems to this list, with each problem's due date specified. Each problem is worth 2 points.

Solutions will be gradually added (and may be hastily written without proofreading).

---

Problem 1 (due Weds 9/4): Let  $A$  be a set and let  $R$  be a relation on  $A$  that is reflexive, symmetric, transitive, and antisymmetric. Prove that for all  $a, a' \in A$  we have  $aRa'$  if and only if  $a = a'$ .

Solution: ( $\Rightarrow$ ): Suppose  $aRa'$ . By symmetry,  $a'Ra$ . By antisymmetry,  $a = a'$ . ( $\Leftarrow$ ): Suppose  $a = a'$ . By reflexivity,  $aRa$ . Since  $a = a'$ , we conclude  $aRa'$ .  $\square$

Problem 2 (due Weds 9/4): Let  $f: A \rightarrow B$  be a function. Suppose that for all  $A_0 \subseteq A$  we have  $f^{-1}(f(A_0)) = A_0$ . Prove that  $f$  is injective.

Solution: Let  $a, a' \in A$  such that  $f(a) = f(a')$ . Then

$$\{a\} = f^{-1}(f(\{a\})) = f^{-1}(\{f(a)\}) = f^{-1}(\{f(a')\}) = f^{-1}(f(\{a'\})) = \{a'\},$$

so  $a = a'$ .  $\square$

Problem 3 (due Weds 9/4): Prove that if a relation is symmetric and antisymmetric, then it is transitive. (So the thing I couldn't think of in class actually doesn't exist, which is nice.)

Solution: Suppose  $R$  is a relation on a set  $A$ , and  $R$  is symmetric and antisymmetric. Let  $a, b, c \in A$  such that  $aRb$  and  $bRc$ . By symmetry,  $cRb$ . By antisymmetry,  $b = c$ . Since  $aRb$  and  $b = c$  we conclude  $aRc$ .  $\square$

---

Problem 4 (due Weds 9/11): Let  $X$  be a set and view the power set  $\mathcal{P}(X)$  as a poset via the partial order  $\subseteq$ . Prove that every subset  $\mathcal{A}$  of  $\mathcal{P}(X)$  has a greatest lower bound. [Added later: If you want to assume  $\mathcal{A}$  is non-empty that's fine, the  $\mathcal{A} = \emptyset$  case is a little weird. (Technically \*every\* element is a lower bound of the empty set, but that's weird.)]

Solution: Let  $\mathcal{A} \subseteq \mathcal{P}(X)$ . Let  $B = \bigcap_{A \in \mathcal{A}} A$ . Then  $B \subseteq A$  for all  $A \in \mathcal{A}$ , so  $B$  is a lower bound of  $\mathcal{A}$ . Also, given any lower bound  $C$  of  $\mathcal{A}$ , we have  $C \subseteq A$  for all  $A \in \mathcal{A}$ , so  $C \subseteq B$ . We conclude that  $B$  is the greatest lower bound of  $\mathcal{A}$ . [Remark: If  $\mathcal{A} = \emptyset$  then, vacuously, every subset of  $X$  is a lower bound of  $\emptyset$ , and so  $X$  itself is the greatest lower bound of  $\emptyset$ . Weird.]  $\square$

Problem 5 (due Weds 9/11): Let  $P$  and  $Q$  be posets with partial orders  $\leq_P$  and  $\leq_Q$ . Let  $\leq_{prod}$  be the product order on  $P \times Q$ . Prove that if every subset of  $P$  has a least upper bound in  $\leq_P$  and every subset of  $Q$  has a least upper bound in  $\leq_Q$ , then every subset of  $P \times Q$  has a least upper bound in  $\leq_{prod}$ .

Solution: Let  $X \subseteq P \times Q$ . Let  $A = \{p \in P \mid (p, q) \in X \text{ for some } q \in Q\}$  and  $B = \{q \in Q \mid (p, q) \in X \text{ for some } p \in P\}$ . Let  $a$  be the least upper bound of  $A$  and let  $b$  be the least upper bound of  $B$ . We claim that  $(a, b)$  is the least upper bound of  $X$ . For any  $(p, q) \in X$  we have  $p \leq_P a$  and  $q \leq_Q b$ , so  $(p, q) \leq_{prod} (a, b)$ , so it is an upper bound. Now let  $(c, d)$  be any upper bound of  $X$ . For any  $(p, q) \in X$  we have  $p \leq_P c$  and  $q \leq_Q d$ . Since  $a$  and  $b$  are the respective least upper bounds,  $a \leq_P c$  and  $b \leq_Q d$ . Thus  $(a, b) \leq_{prod} (c, d)$ , and we conclude that  $(a, b)$  is the least upper bound of  $X$ .  $\square$

Problem 6 (due Weds 9/11): Consider the Cantor set  $C = \prod_{\mathbb{N}} \{0, 1\}$ , so elements of  $C$  are infinite binary sequences  $(a_1, a_2, \dots)$ . Let  $\psi: C \rightarrow \mathbb{R}$  be the function sending  $(a_1, a_2, \dots)$  to  $\sum_{i=1}^{\infty} \frac{a_i}{10^i}$  (don't worry, this infinite series converges thanks to some calculus thing, so this really is an element of  $\mathbb{R}$ ). Prove that  $\psi$  is injective.

Solution: Let  $\vec{a} = (a_1, a_2, \dots), \vec{b} = (b_1, b_2, \dots) \in C$  such that  $\psi(\vec{a}) = \psi(\vec{b})$ , so  $\sum_{i=1}^{\infty} \frac{a_i}{10^i}$  equals  $\sum_{i=1}^{\infty} \frac{b_i}{10^i}$ . By calculus stuff,  $\sum_{i=1}^{\infty} \frac{a_i - b_i}{10^i} = 0$ . Multiplying both sides by 10 we get  $a_1 - b_1 + \sum_{i=1}^{\infty} \frac{a_{i+1} - b_{i+1}}{10^i} = 0$ . Since all the  $a_i$  and  $b_i$  are 0 or 1, we know that  $\sum_{i=1}^{\infty} \frac{a_{i+1} - b_{i+1}}{10^i}$  is bounded above and below by  $\sum_{i=1}^{\infty} \frac{1}{10^i} = 1/9$  and  $\sum_{i=1}^{\infty} \frac{-1}{10^i} = -1/9$ . Hence  $|a_1 - b_1| \leq 1/9$ , so  $a_1 - b_1 = 0$ . Now multiplying our new equation  $\sum_{i=1}^{\infty} \frac{a_{i+1} - b_{i+1}}{10^i} = 0$  on both sides by 10, we get  $a_2 - b_2 + \sum_{i=1}^{\infty} \frac{a_{i+2} - b_{i+2}}{10^i} = 0$ , and the same argument shows  $a_2 - b_2 = 0$ . Repeating this, we get  $a_i - b_i = 0$  for all  $i$ , and conclude that  $\vec{a} = \vec{b}$ .  $\square$

Problem 7 (due Weds 9/18): Let  $X = \{a, b\}$  and let  $\mathcal{T} \subseteq \mathcal{P}(X)$  satisfy  $\emptyset, X \in \mathcal{T}$ . Prove that  $\mathcal{T}$  is a topology.

Solution: Since  $X$  is finite and  $\mathcal{T}$  contains  $\emptyset$  and  $X$ , it suffices to show that for any  $U, V \in \mathcal{T}$  we have  $U \cup V, U \cap V \in \mathcal{T}$ . If either  $U$  or  $V$  is  $\emptyset$  or  $X$ , or if  $U = V$ , this is trivially true. Now assume none of these are the case, so  $U$  and  $V$  each have exactly one element, and they are different. Without loss of generality  $U = \{a\}$  and  $V = \{b\}$ , and now the result is clear.  $\square$

Problem 8 (due Weds 9/18): Let  $X$  be a set and fix  $x_0 \in X$ . Let  $\mathcal{T}_{x_0} = \{U \subseteq X \mid x_0 \in U\} \cup \{\emptyset\}$ . Prove that  $\mathcal{T}_{x_0}$  is a topology.

Solution: We are given that  $\emptyset \in \mathcal{T}_{x_0}$ , and clearly  $X \in \mathcal{T}_{x_0}$  since  $x_0 \in X$ . Now let  $U_\alpha \in \mathcal{T}_{x_0}$  for all  $\alpha \in \Lambda$ . If all  $U_\alpha$  are empty, then their union is empty, hence open. If some  $U_\alpha$  is non-empty, it must contain  $x_0$ , so the union of all of them contains  $x_0$ , and hence is open. Finally, let  $U, V \in \mathcal{T}_{x_0}$ . If  $U$  or  $V$  is empty, so is their intersection, hence  $U \cap V$  is open. If neither is empty, they both contain  $x_0$ , hence so does their intersection, so it is open.  $\square$

Problem 9 (due Weds 9/18): Let  $X = \mathbb{N}$  with the finite complement topology. Let  $\mathcal{B} = \{U \subseteq X \mid 2024 \leq |X \setminus U| < \infty\}$ . Prove that  $\mathcal{B}$  is a basis for a topology, and the topology it generates equals the finite complement topology.

Solution: Let  $x \in X$ . Since  $X$  is infinite, there exists  $S \subseteq X \setminus \{x\}$  with  $|S| = 2024$ . Set  $U = X \setminus S$ , so  $X \setminus U = S$ . Now  $U \in \mathcal{B}$ , and  $x \in U$ , thus proving the first basis axiom. Next let  $x \in U \cap V$  for  $U, V \in \mathcal{B}$ . Note that  $X \setminus (U \cap V) = (X \setminus U) \cup (X \setminus V)$ . Since  $2024 \leq |X \setminus U|, |X \setminus V| < \infty$ , we have  $2024 \leq |(X \setminus U) \cup (X \setminus V)| < \infty$ , so in fact  $U \cap V \in \mathcal{B}$ . This proves the second axiom. Now we must show that the topology generated by  $\mathcal{B}$  equals the finite complement topology. Every set in  $\mathcal{B}$  has finite complement, so one direction is clear. Now let  $x \in W$  for an arbitrary  $W \subseteq X$  with  $|X \setminus W| < \infty$ . Since  $X$  is infinite, we can choose  $S \subseteq X \setminus W$  with  $|S| = 2024$ . Set  $W' := W \setminus S$ . Since  $x \in W$  we have  $x \notin S$ , so  $x \in W'$ . Also,  $|X \setminus W'| = |X \setminus (W \setminus S)| \geq |S| = 2024$ , so  $W' \in \mathcal{B}$ . Now  $x \in W' \subseteq W$  reveals that every  $W$  open in the finite complement topology is also open in the topology generated by  $\mathcal{B}$ , and we are done.  $\square$

---

No homework due 9/25, just the exam.

---

Problem 10 (due Weds 10/2): Let  $X = \mathbb{N}$  with the finite complement topology and let  $S \subseteq X$ . Prove that the interior  $\overset{\circ}{S}$  is empty if and only if  $X \setminus S$  is infinite.

Solution: ( $\Rightarrow$ ): Contrapositively, suppose that  $X \setminus S$  is finite. Then  $S$  is non-empty and open, and so  $\overset{\circ}{S} = S$  is non-empty. ( $\Leftarrow$ ): Contrapositively, suppose  $\overset{\circ}{S} \neq \emptyset$ , so there exists a non-empty open  $U$  contained in  $S$ . Thus  $X \setminus U$  is finite, and hence so is its subset  $X \setminus S$ .  $\square$

Problem 11 (due Weds 10/2): Let  $X$  be a topological space in which every subset is either closed or dense. Prove that every subset of  $X$  containing a non-empty open set is open.

Solution: Let  $\emptyset \neq U \subseteq S \subseteq X$  for  $U$  open in  $X$ . We claim that  $S$  is open. Since  $X \setminus S$  is contained in the closed set  $X \setminus U \neq X$ , the closure of  $X \setminus S$  is too, and hence cannot equal  $X$ . This means  $X \setminus S$  is not dense, so it must be closed, and thus  $S$  is open.  $\square$

Problem 12 (due Weds 10/2): Let  $X$  and  $Y$  be [added: non-empty] [oh shoot also not singletons!] sets with the finite complement topology. Prove that if the product topology on  $X \times Y$  equals the finite complement topology on  $X \times Y$ , then  $X$  and  $Y$  are both finite.

Solution: Suppose  $X$  and  $Y$  are not both finite, so without loss of generality  $X$  is infinite. Fix  $y_0 \in Y$  (here we needed to assume the sets are non-empty). Now  $Y \setminus \{y_0\}$  is open in  $Y$ , so  $X \times (Y \setminus \{y_0\})$  is open in the product topology on  $X \times Y$ . But the complement of  $X \times (Y \setminus \{y_0\})$  in  $X \times Y$  is  $X \times \{y_0\}$ , which is infinite [and not all of  $X \times Y$  since we rule out  $X$  and  $Y$  being singletons], so  $X \times (Y \setminus \{y_0\})$  is not open in the finite complement topology.  $\square$

---

Problem 13 (due Weds 10/9): Let  $f: X \rightarrow Y$  be a continuous surjection of topological spaces. Suppose that  $Y$  is Hausdorff, and that for all  $y \in Y$  the subspace  $f^{-1}(\{y\})$  of  $X$  is open and Hausdorff. Prove that  $X$  is Hausdorff.

Solution: Let  $x \neq x'$  in  $X$ . First suppose  $f(x) \neq f(x')$ . Since  $Y$  is Hausdorff we can choose disjoint open neighborhoods  $V$  and  $V'$  of  $f(x)$  and  $f(x')$  in  $Y$ . Let  $U = f^{-1}(V)$  and  $U' = f^{-1}(V')$ . Now  $x \in U$ ,  $x' \in U'$ ,  $U$  and  $U'$  are open in  $X$ , and they are disjoint. Alternately, suppose  $f(x) = f(x')$ , call it  $y$ , so  $x, x' \in f^{-1}(\{y\})$ . Since  $f^{-1}(\{y\})$  is Hausdorff, we can choose disjoint open neighborhoods of  $x$  and  $x'$  in  $f^{-1}(\{y\})$ , and since  $f^{-1}(\{y\})$  is open in  $X$ , these neighborhoods are open in  $X$ , so we are done.  $\square$

Problem 14 (due Weds 10/9): Prove that  $C = \prod_{\mathbb{N}}\{0, 1\}$  (with the product topology) is Hausdorff.

Solution: Let  $(a_1, a_2, \dots) \neq (b_1, b_2, \dots)$  in  $C$ , so these are infinite sequences of 0s and 1s. Choose some  $i \in \mathbb{N}$  such that  $a_i \neq b_i$ , say without loss of generality that  $a_i = 0$  and  $b_i = 1$ . Let  $U_i = \{0\}$ ,  $V_i = \{1\}$ , and  $U_j = V_j = \{0, 1\}$  for all  $j \neq i$ . Now  $U_1 \times U_2 \times \dots$  and  $V_1 \times V_2 \times \dots$  are open neighborhoods of  $(a_1, a_2, \dots)$  and  $(b_1, b_2, \dots)$  respectively, and they are disjoint since the  $i$ th entry of any element of their intersection would have to be simultaneously 0 and 1.  $\square$

Problem 15 (due Weds 10/9): Let  $X$  be a space with the finite complement topology. Prove that if  $X$  is Hausdorff then it is finite.

Solution: Suppose  $X$  is Hausdorff. Let  $x \neq y$  in  $X$ . Choose disjoint open neighborhoods  $U$  of  $x$  and  $V$  of  $y$  in  $X$ . Since  $U$  and  $V$  are non-empty and open, they have finite complement. Now  $U \cap V = \emptyset$  implies  $(X \setminus U) \cup (X \setminus V) = X$ , so  $X$  is a union of two finite sets, hence is finite.  $\square$

---

No homework over October break (nothing due 10/16).

---

Problem 16 (due Weds 10/23): Let  $(X, d)$  be a metric space and let  $x \in X$ . Let  $f_x: X \rightarrow \mathbb{R}$  be  $f_x(y) := d(x, y)$ . Prove that  $f_x$  is continuous.

Solution: Let  $(a, b)$  be a basic open set in  $\mathbb{R}$ , so  $a < b$ . Then  $f_x^{-1}(a, b) = \{y \in X \mid a < d(x, y) < b\}$ . This is equal to  $\{y \in X \mid y < b\} \cap \{y \in X \mid a < y\}$ . That first set is the basic open set  $B_b(x)$ , so it suffices to show that  $\{y \in X \mid a < y\}$  is open. Indeed, for any  $y$  in this set, the open ball centered at  $y$  with radius  $(y - a)/2$  is fully contained in this set.  $\square$

Problem 17 (due Weds 10/23): Let  $(X, d)$  be a metric space. Let  $Y \subseteq X$ . The *induced metric*  $d_Y$  on  $Y$  is defined to be  $d_Y(y, y') := d(y, y')$ . (This is clearly a metric, you don't have to prove that.) Prove that the metric topology on  $Y$  coming from  $d_Y$  equals the subspace topology on  $Y$  coming from  $X$ .

Solution: First let us prove that every basic open set in the metric topology is open in the subspace topology. A basic open set in the metric topology is of the form  $\{y' \in Y \mid d_Y(y, y') < r\}$  for some  $y \in Y$  and  $r \in \mathbb{R}$ . This equals  $Y \cap \{x' \in X \mid d(y, x') < r\}$ , i.e.,  $Y$  intersect a (basic) open set in  $X$ , so indeed this is open in the subspace topology. Next let us prove that every open set in the subspace topology is open in the metric topology. Let  $Y \cap U$  be an arbitrary open set in the subspace topology, so  $U$  is open in  $X$ . Let  $y \in Y \cap U$ . Choose  $r \in \mathbb{R}$  such that  $B_r(y) \subseteq U$  (here  $B_r(y)$  is the open ball of radius  $r$  centered at  $y$  in  $X$ ). Now  $Y \cap B_r(y) = \{y' \in Y \mid d(y, y') < r\} = \{y' \in Y \mid d_Y(y, y') < r\}$ , which is a basic open set in the metric topology. Hence  $y \in Y \cap B_r(y) \subseteq Y \cap U$  shows that  $Y \cap U$  is open in the metric topology.  $\square$

Problem 18 (due Weds 10/23): Let  $X$  be a set. A function  $d: X \times X \rightarrow \mathbb{R}$  is a *pseudometric* if it satisfies the same axioms as a metric, except we allow  $d(x, y) = 0$  even if  $x \neq y$ . Define  $B_r(x) = \{y \in X \mid d(x, y) < r\}$  for  $r \in \mathbb{R}$ . It turns out the collection of all  $B_r(x)$  forms a basis for a topology on  $X$  (you don't need to prove this). Prove that if  $X$  with this pseudometric topology coming from  $d$  is Hausdorff, then  $d$  is actually a metric, i.e.,  $d(x, y) = 0$  implies  $x = y$ .

Solution: Suppose the pseudometric topology on  $X$  is Hausdorff, and suppose  $x, y \in X$  with  $d(x, y) = 0$ . We must show that  $x = y$ . Since  $X$  is Hausdorff, it suffices to show that every (basic) open neighborhood of  $x$  intersects every (basic) open neighborhood of  $y$ . Let  $x \in B_r(z)$  and  $y \in B_s(w)$ . Since  $x \in B_r(z)$  we know  $d(z, x) < r$ , and since  $d(x, y) = 0$  we get  $d(z, y) \leq d(z, x) + d(x, y) < r$ , so  $y \in B_r(z)$ . Thus  $y \in B_r(z) \cap B_s(w)$ , so this is non-empty.  $\square$

---

No homework due 10/30, just the exam.

---

Problem 19 (due Weds 11/6): Let  $X = \mathbb{Z}$  with the “particular point” topology, where a non-empty set is open if and only if it contains 0. Prove that  $X$  is path connected.

Problem 20 (due Weds 11/6): Prove that  $\mathbb{R}$  with the finite complement topology is path connected. [As a remark, it turns out  $\mathbb{Z}$  with the finite complement topology is not path connected, but this is kind of hard to prove.]

Problem 21 (due Weds 11/6): Let  $(X, d)$  be a metric space. A *geodesic* is a path  $p: [a, b] \rightarrow X$  such that for all  $t, t' \in [a, b]$  we have  $d(p(t), p(t')) = |t - t'|$ . Call  $X$  a *geodesic space* if for all  $x, y \in X$  there exists a geodesic from  $x$  to  $y$ . Prove that any geodesic space is locally path connected.

---